



Mirror of Research in Veterinary Sciences and Animals (MRVSA)

Journal homepage: <http://mrvisa.com/>

E-ISSN 2307-8073; 2520-324X (Print)



Crossref



DOAJ

DIRECTORY OF
OPEN ACCESS
JOURNALS

Detection of bovine viral diarrhea-mucosal disease (BVD-MD) virus in Dromedary camel in Iraq using ELISA/ A preliminary study

Khawlah Moh Imran Al-Rubayie ^{1*}¹ Department of internal and preventive Medicine/ College of Veterinary Medicine/ Baghdad university- Baghdad –Iraq**ARTICLE INFO****Received:** 05.04.2016**Revised:** 10. 06.2016**Accepted:** 12.06.2016**Publish online:** 25.06.2016***Corresponding author:**husein.abdulridha@uokufa.edu.iq**Abstract**

Bovine Viral Diarrhea-Mucosal Disease (BVD- MD) virus causes a high economic losses. BVD-MD infects a wide range of domestic animals (cattle, buffaloes, sheep and goat). The causative agent,

bovine viral diarrhea virus (BVDV) is a member of the Pestivirus genus of the family Flaviviridae that composed of two genotypes (BVDV-1, BVDV-2) and each genotype has 2 biotypes: non-cytopathic (NCP) and cytopathic (CP). Only NCP strains of BVDV produce Persistently Infected (PI) animals. Eighty- eight blood samples were collected from dromedary camels in different areas surrounding Baghdad (Al-Shula, Abu- grab and Al-Fudhailiyih), as well as from Karbala, Najaf and Babylon governorates. All serum samples were examined using a BVDV specific indirect enzyme – linked –immunosorbent assay (ELISA) specific to BVD-MD virus antibody. Totally, 12 out of 88 (13.63%) serum samples revealed positive reaction for (BVD-MD) virus antibodies. Moreover, 5 / 22 (5.68%), 2 /22(2.27%), 4/22 (4.54%) and 1/22 (1.13%) revealed positive reaction from Baghdad, Karbala, Najaf and Babylon respectively. No significant differences were seen between male and female. In conclusion, this preliminary study approved presences of positive camels for BVD-MD virus. The author recommends to do another future study that included large numbers of camels and from different areas in Iraq, as well as observe the congenital anomalies and abortion in newly borne camels.

To cite this article: Khawlah Moh Imran Al-Rubayie. (2016). Detection of bovine viral diarrhea-mucosal disease (BVD-MD) virus in Dromedary camel in Iraq using ELISA/ preliminary study. 70-74. DOI: [10.22428/mrvsa.2307-8073.2014.002180.x](https://doi.org/10.22428/mrvsa.2307-8073.2014.002180.x)

Keywords: ELISA, BVD-MD virus, Camel, Abu- grab.

Introduction

Bovine viral diarrhea virus (BVDV) is a member of the genus Pestivirus in family Flaviviridae (Franki *et al.*, 1991; Pringle, 1999). Each of the two genotypes has two biotypes, non-cytopathic (NCP) and cytopathic (CP) (Peterhans *et al.*, 2010). The (NCP) genotype is series than (CP) genotype. (Amstel and Kennedy, 2010). The BVDV, border

disease virus (BDV) of sheep and classical swine fever virus (CSFV) are antigenically related, (Nettleton *et al.*, 1998). Bovine viral diarrhea (BVD) is a worldwide distribution infectious disease of cattle (Nettleton, 1995).

BVDV infections involve mainly respiratory, enteric and reproductive organs accompanied with increased risk of retained placenta and clinical mastitis, (Niskanin *et al.*, 1995). BVDV caused intrauterine death, stillbirth, and weak calf syndrome with congenital deformities, neonatal respiratory disorders and acute hemorrhagic gastroenteritis in adult dromedaries. Acute infection induced immunosuppression that potentiate secondary bacterial and viral disease of the respiratory and enteric tract of persistently infected (PI) animals. The (PI) animal remain lifelong shedding large quantities of virus in secretion and excretions (carriers) (Alenius *et al.*, 1996). BVDV is a significant economic disease of ruminant, which is endemic in the majority of countries throughout the world. Review of literature, revealed scarce information regarding BVD in camel in Iraq. So this a preliminary study was designed to detect the infected camels with BVDV-MD in Iraq serologically

Material and methods

Eighty eight Blood samples were collected from dromedary camels from slaughter houses around Baghdad city (Abu-grab, Al-Shula, Al-Fudaiyia) as well as Karbala, Najaf and Babylon governorates. Blood samples were collected in tubes without anticoagulant and kept in cold box and transfer to the laboratory for further processing. Serum samples were separated from each samples and stored at (-20 °C) until used.

Serological test

Elisa Kits (antibody ELISA kits) were purchased from Belgium BIO-X diagnostics. The ELISA procedures were done according the instructions of the manufacture.

Results and discussion

ELISA antibody test were done for (88) serum samples. Totally, 12 (13.63%). out of 88 samples revealed positive results for BVDV antibodies (Table. 1).

Moreover, 5 / 22 (5.68%), 2 / 22 (2.27%), 4/22 (4.54%) and 1/22 (1.13%) revealed positive reaction from Baghdad, Karbala, Najaf and Babylon respectively (Table.1). Meanwhile, Baghdad governorate revealed the high number of positive camels and 5 out of the 22 samples revealed positive reaction. The positive samples were 1/7, 3/8 and 1/7 from Al-Shula, Abu-grab and Al- Fudhailiyih respectively (Table. 2). According to the sex, the samples were collected from 30 and 58 male and female respectively. Moreover, there were 4 (13.3%) and 8 (13.7%) positive samples from male and female respectively (Table.3). Statistical analysis revealed non-significant difference between male and female.

Table.1: Shows the number and percentage of positive samples in different governorate

Name of governorate	No. Sample	Positive	Percentage of infection
Baghdad	22	5	5.68%
Karbala	22	2	2.27%
Najaf	22	4	4.54%
Babylon	22	1	1.13.%
Total	88	12	13.63 %

Table.2: Shows the number and percentage of positive samples in different area surrounding Baghdad governorate

No. of examined samples	Name of area	No. of animals	Positive	Percentage
22	Al-Shula	7	1	4.54%
	Abu-grab	8	3	13,63%
	Al-Fudhailiyih	7	1	4,54%
Total		22	5	22.72 %

Table .3: Shows the number of positive samples in male and female with percentage

No. of examined animals	Sex	No.	Positive	Percentage
88	Male	30	4	13.3%
	Female	58	8	13.7%

BVD virus was detected in cattle in Iraq (Al Rodhan 2005). It was also approved in buffaloes ((Al-Rubayie and Hasso, 2012) using ELISA antigen and antibody. The results revealed presence of antibody against BVDV. This result is in agreement with (Doyle and Heuschele, 1983). The percentage of infection in Baghdad was (22.72 %). This result is compatible previous studies in Dromedary (18%) in Saudi Arabia (Al-Afaleq *et al.* 2006).

However, the results of this study is disagreed with Taha, (2007) in UAE dromedary that approved the sero-prevalence (negative) (Taha, 2007). The results of the current study revealed high percentage of positive animals in compare with previous studies. This might occurred due to mixed breeding management between animals in Baghdad especially the animals that act as a source of BVDV infection in camelids. The author believe that the infection occurred via the oronasal mucos, which occurs most probably after inhalation of viral particles which are present in body fluids of infected animals (Byers *et al.*, 2011). In conclusion this study approved that the antibody of (BVD–MD)

Khawlah Moh Imran Al-Rubayie ., (2016); 5 (Special issue), 70-74.

1st Iraqi colloquium on camel diseases and management.

Mirror of Research in Veterinary Sciences and Animals

virus was detected by specific (BVDV) antibody ELISA kits in dromedary camel. This result indicates the presence of the disease in Iraq in dromedary. The author recommends to do another future study that included large numbers of camels from different areas in Iraq. Polymerase chain reaction (PCR) is recommended with Antigen-enzyme-linked immunosorbent assay, and also skin biopsy with immunohistochemistry, (IHC) – antigen detection. Virus isolation and typing, sequencing are necessary also, in addition to observe the congenital anomalies and abortion in newly borne camels.

References

Al-Afaleq A, Abu-Elzein E M E. and Hegazy A A. (2006). Sero-surveillance for antibodies against some viral diseases of livestock in camels (*Camelus dromedarius*) in Saudi Arabia. Proceedings of the International Scientific Conference on Camels, (ISCC' 06), Qassim University, Kingdom of Saudi Arabia, pp: 338-346.

Amstel S V and Kennedy M. (2010). Bovine viral diarrhea infections in new world camelids-A review. Small Ruminant Res. 91: 121-126. DOI: 10.1016/j.smallrumres.2010.03.018

Anonymous (2006) a. Bovine viral diarrhea virus in camelids. Proceedings of the International Camelid Health Conference for Veterinarians, (ICHCV' 06), Ohio State University, USA. pp: 275-278.

Al-Rodhan, MA. (2005). Bovine viral diarrhea virus isolation and partial Characterization. PhD Thesis, Coll. Vet. Med. Uni. Bagh.

Byers SR, Evermann JF, Bradway DS, Grimm A L. and Ridpath J F et al., (2011). The effects of exposure of susceptible alpacas to alpacas persistently infected with bovine viral diarrhea virus. Can. Vet. J. 52: 263-271.

Carman S, Carr N, DeLay J, Baxi M. and Deregt D et al., (2005). Bovine viral diarrhea virus in alpaca: Abortion and persistent infection. J. Vet. Diag. Invest., 17: 589-593. PMID: 16475521

Doyle L G and Heuschele W P. (1983). Bovine viral diarrhea virus infection in captive exotic ruminants. JAVMA, 183: 1257-1259. PMID: 6315662.

Franki RIB, Fauquet CM, Knudson DL, Brown F (1991). Classification and nomenclature of viruses; fifth report of the international committee on taxonomy of viruses. Arch Virol. 223–233.

Hegazy A A, Fahmy L S, Saber M S, Aboellail TA and Yousif A A et al., (1998). Bovine virus diarrhea infection causes reproductive failure and neonatal mortality in the dromedary camel. Proceedings of the International Meeting on Camel Production and Future Perspectives. Al Ain, UAE, 2-3.

Khawlah Moh Imran Al-Rubayie ., (2016); 5 (Special issue), 70-74.

1st Iraqi colloquium on camel diseases and management.

Mirror of Research in Veterinary Sciences and Animals

Hegazy A A, Lotfia S F and Saber M S. (1993). Prevalence of antibodies common in viral diseases of domestic animals among camels in Egypt.

Kapil S, Yeary T and Evermann J F. (2009). Viral diseases of new world camelids. Vet. Clin. Food Anim., 25: 323-337. PMID: 19460643

Kelling C. (2008). Prevalence of BVDV infected US alpaca herds and factors associated with BVDV seropositive herd status. Proceedings International Camelid Health Conference for Veterinarians, (ICHCV' 08), Ohio State University, USA. pp. 176-181.

Khawlah M.I. Al-Rubayie, Saleem A. Hasso (2012). Detection of bovine viral diarrhoea mucosal disease BVD-Din buffaloes and cows using ELISA, the Iraqi J.Vet.Med.361:45-50.

Mudry M, Meylan M, Regula G, Steiner A and Zanoni R *et al.*, (2010). Epidemiological study of pestiviruses in South American camelids in Switzerland. J. Vet. Intern. Med., 24: 1218-1223. DOI: 10.1111/j.1939-1676.2010.0577.x

Nettleton P F, Gilray J A, Russo P, Dlissi E. (1998). Border disease of sheep and goats. Veterinary Research, 29, 327 340.

Niskanen R. (1993). Relationship between the levels of antibodies to bovine viral-diarrhoea virus in bulk tank milk and the prevalence of cows exposed to the virus. Veterinary Record. 133: 341-344.

Nettleton P F & Entrican G. (1995). Ruminant pestivirus: A review. Br.Vet. Jour. 151, 615-641.

Peterhans E, Bachofen C, Stalder H and Schweizer M (2010). Cytopathic Bovine Viral Diarrhoea Viruses (BVDV). Emerging pestiviruses doomed to extinction. Vet. Res. 41: 1-14. DOI: 10.1051/vetres/2010016

Taha T H. (2007). Pathogens affecting the reproductive system of camels in the United Arab Emirates. M. Sc., Thesis, Swedish University of Agricultural Sciences, Uppsala.

Wernery U. (2012). Bovine viral diarrhoea-an emerging disease in camelids a review.